

DISCRETE SPHERES AND ARITHMETIC PROGRESSIONS IN PRODUCT SETS

ABSTRACT. We prove that if B is a set of N positive integers such that $B \cdot B$ contains an arithmetic progression of length M then $N \geq \pi(M) + M^{2/3-o(1)}$. On the other hand, there are examples for which $N < \pi(M) + M^{2/3}$. This improves previously known bounds of the form $N = \Omega(\pi(M))$ and $N = O(\pi(M))$, respectively.

The main new tool is a reduction of the original problem to the question of an approximate additive decomposition of the 3-sphere in $\mathbb{F}_3^n$ which is the set of 0-1 vectors with exactly three non-zero coordinates. Namely, we prove that such a set cannot be contained in a sumset $A + A$ unless $|A| \gg n^2$.

1. INTRODUCTION

One of the main objectives of additive combinatorics is to show that a set cannot be simultaneously additively and multiplicatively structured. In particular, if one defines the sumset

$$A + A = \{a + a' \mid a, a' \in A\}$$

and the product set

$$A \cdot A = \{aa' \mid a, a' \in A\}$$

of a set A , the sum-product conjecture of Erdős and Szemerédi claims that the size of at least one of these sets is substantially larger than that of the original set. Originally, the conjecture was formulated only for integer sets, but later it became clear that conceivably it should hold in a much more general setting, so now it is usually stated for $A \subset \mathbb{C}$. More precisely,

$$\max\{|A \cdot A|, |A + A|\} \geq C_\epsilon |A|^{2-\epsilon}, \quad (1)$$

where $\epsilon > 0$ can be taken arbitrarily small and C_ϵ is some constant which depends only on ϵ . There are numerous results towards this conjecture in different flavours but it is still far from resolution. We refer the interested reader to [3] for an overview of this topic¹.

On the other hand, very little is known about sets which are themselves of the form $B \cdot B$ or $B + B$, or subsets thereof. Clearly, such sets must be as well

Date: October 20, 2015.

2000 Mathematics Subject Classification. 11B25 (primary).

Key words and phrases. product sets, arithmetic progressions.

¹Further progress has been made since the paper [3] was published, but it still gives a great overview of the sum-product phenomenon. See also the standard reference [4].

multiplicatively resp. additively structured in a certain sense, but perhaps in a more subtle way. For example, it is an open problem to algorithmically decide whether a given set A is of the form $B + B$, see Problem 4.11 in [1].

By analogy to the classical sum-product phenomenon, one might expect that $B \cdot B$ (resp. $B + B$) cannot contain large additively (resp. multiplicatively) structured pieces, with some freedom left in defining what the actual structure is. Some results in this direction have been obtained by Roche-Newton and the author [2], [5], where the structure is defined in terms of the additive (resp. multiplicative) energy. In the extreme case of a product set $B \cdot B$ containing an arithmetic progression the author has shown [6] that the maximal length of the arithmetic progression must be $O(|B| \log |B|)$, at least in the integer case, together with an example exhibiting a progression of length $\Omega(|B| \log |B|)$. The goal of the present note is to further refine these bounds by recasting the original problem to $\mathbb{F}_3^n$ and then applying linear algebra and additive combinatorics. This allows one to surpass the previous bound which was based mainly on number theoretic arguments. Finally, Lemma 4 below about additive decomposition of discrete spheres in $\mathbb{F}_3^n$ might be of independent interest.

Throughout the paper we use the standard notation $X \ll Y$, $Y \gg X$, $X = O(Y)$, $Y = \Omega(X)$ which means that there is a constant C independent of X and Y such that $|X| \leq CY$. The asymptotic little-oh notation $o(1)$ denotes a quantity which becomes arbitrarily small if N (or, equivalently, M) is sufficiently large.

2. MAIN RESULT

Theorem 1. *If B is a set of N positive integers, such that $B \cdot B$ contains an arithmetic progression of length M then $N \geq \pi(M) + M^{2/3-o(1)}$. Moreover we can construct examples for which $N < \pi(M) + M^{2/3}$.*

Obviously, a lower (upper) bound on N in terms of M is equivalent to an upper (lower) bound on M in terms of N . We will prove Theorem 1 by bounding M in terms of N rather than what is stated, as it is slightly more convenient for calculations.

Proof of lower bound on M . We will construct B so that $\{1, \dots, M\} \subset B \cdot B$ for M as large as possible. Let $B = \{n \leq x\} \cup \{p \leq M : p \text{ prime}\}$, where $x = M^{2/3}$. For $m \leq M$ let p be the largest prime factor of m ; if $p \geq y := M/x$ then $m/p \leq M/y = x$ and so $p, m/p \in B$. If $m \leq x$ then $m, 1 \in B$. If $p < y$ and $x < m \leq M$ then let d be the largest divisor of m that is $\leq x$ so that $x/y < d \leq x$ and $m/d < M/(x/y) = M^2/x^2 = x$ so that $d, m/d \leq x$. Thus, both $d, m/d$ are in B and $m \in B \cdot B$. Here $|B| = \pi(M) + M^{2/3} - \pi(M^{2/3})$. $\square$

Lemma 1. *Suppose that there exist a set B of N positive integers, such that there is an M -term arithmetic progression $A = \{a + md : 1 \leq m \leq M\} \subset B \cdot B$. Then*

there exist sets such that if $g = (a, d)$ with $a = gu$ and $d = gv$ then $(v, g) = 1$. For these sets we have the following property:

If $\mathcal{M} \subset \{m : 1 \leq m \leq M\}$ is such that for each $m \in \mathcal{M}$ there exists a prime p_m for which $p_m | u + vm$ and $p_m \nmid u + vm'$ for all $m' \in \mathcal{M} \setminus \{m\}$, then $|B| \geq |\mathcal{M}|$.

Proof. Choose such sets A, B with the product of the elements of B minimal. If $(v, g) \neq 1$ then there exists a prime p and exponents $e > f \geq 1$ with $p^e || d$ and $p^f || a$. Therefore $p^f ||$ every element of A , which means that if that element equals $b_1 b_2$ then

$$\text{either } 0 < v_p(b_1), v_p(b_2) < f; \quad \text{or } v_p(b_1) = 0 \text{ and } v_p(b_2) = f,$$

(where $v_p(b)$ is the exact power of p dividing b).

If $f = 1$ then let $A' = \{a/p : a \in A\}$ and

$$B' := \{b \in B : v_p(b) = 0\} \cup \{b/p : b \in B \text{ \& } 0 < v_p(b) = f\}.$$

Otherwise, let $A' = \{a/p^2 : a \in A\}$ and

$$B' := \{b \in B : v_p(b) = 0\} \cup \{b/p : b \in B \text{ \& } 0 < v_p(b) < f\} \cup \{b/p^2 : b \in B \text{ \& } v_p(b) \geq f\}.$$

One easily verifies that $A' \subset B' \cdot B'$ with $|A'| = M$, $|B'| = N$ and the product of the elements of B' is much smaller than the product of the elements of B . This contradicts our assumption of minimality.

Now we prove the second part of the lemma. Let $\mathcal{P}$ be the set of primes p_m with $m \in \mathcal{M}$, so $|\mathcal{P}| = |\mathcal{M}| := M_p$. Define a map $\rho : \mathbb{Z} \rightarrow (\mathbb{F}_q)^{M_p}$, where q is some prime number defined in due course, as follows. If $x = D \prod_{m \in \mathcal{M}} p_m^{e_m}$, then $\rho(x)_i = e_i \pmod q$. In words, the i th coordinate of $\rho(x)$ is $v_{p_i}(x)$ modulo q . Clearly, $\rho(A) \subset \rho(B) + \rho(B)$ and $|B| \geq |\rho(B)|$.

Now let $\rho_g = \rho(g)$ and define $B' = \rho(B) - \rho(g)/2$, where the division is coordinate-wise. It is easy to see that $\rho(u + vm) \in B' + B'$ for any $m \in \mathcal{M}$. Moreover, by the hypothesis of the lemma we can choose q such that $\rho(u + vm)$ has only one non-zero coordinate, namely m . But then obviously the dimension of $\text{Span}(B')$ must be M_p , so $|B'| \geq M_p = |\mathcal{M}|$. The desired property now follows. $\square$

Lemma 2. *Using the notation of Lemma 1, if $|B| \leq \pi(M) + M^{2/3}$, one can assume that $v = 1$, $u \ll M^{2/3} \log M$.*

Proof of upper bound on M . We begin by using the lemma. If p is a prime $\geq M$ then it can divide at most one $u + mv$ (since $(u, v) = 1$). Let

$$\mathcal{M} := \{m : 1 \leq m \leq M \text{ \& } \exists p \geq M : p | u + mv\}.$$

Now if $1 \leq m \leq M$ and $m \notin \mathcal{M}$ then all prime factors of $u + mv$ are $< M$. For each prime $p < M$ select m_p for which $v_p(u + m_p v)$ is maximized and then

$\mathcal{E} := \cup_{p < M} \{m_p\}$. We claim that

$$\prod_{\substack{1 \leq m \leq M \\ m \notin \mathcal{M} \cup \mathcal{E}}} (u + mv) \text{ divides } (M-1)!$$

Indeed, the left hand side has only prime divisors less than M and for such a prime $p < M$ the maximal power of p which divides the left hand side is at most

$$\sum_{k=1}^{\infty} \left[\frac{M-1}{p^k} \right].$$

This is because if $k \leq v_p(u + m_p v)$ the only elements of $\{u + mv\}$ divisible by p^k are of the form $u + (m_p + jp^k)v, j \in \mathbb{Z}$. But the same total power of p also divides $(M-1)!$.

Let $L := |\mathcal{M} \cup \mathcal{E}| \leq |\mathcal{M}| + |\mathcal{E}| < |B| + \pi(M-1)$. Hence

$$\prod_{m=1}^{M-L} (u + mv) \leq (M-1)! \quad (2)$$

If we assume that $|B| \leq \{1 + o(1)\}M/\log M$ then $L \leq \{2 + o(1)\}M/\log M$ by the prime number theorem. If $v \geq 9$ then (2) yields², using Stirling's formula,

$$\begin{aligned} (9/e^{2+o(1)})^M (M/e)^M &\ll (9(M-L)/e)^{M-L} \\ &\ll (M-L)! v^{M-L} \leq (M-1)! \ll (M/e)^M, \end{aligned}$$

which gives a contradiction for M sufficiently large.

If $u \geq 7M$ then (2) yields, using Stirling's formula,

$$\begin{aligned} (8^8/e^{2+o(1)})^M (M/e)^{8M} &\ll ((8M-L)/e)^{8M-L} \\ &\ll (8M-L)! \leq (7M)!(M-1)! \ll (7M/e)^{7M} (M/e)^M, \end{aligned}$$

which gives a contradiction for M sufficiently large.

Hence we may assume that $v \leq 9$ and $u < 7M$, so that each $u + mv \leq 16M$. Any element of the arithmetic progression $\{u + mv\}$ is divisible by at most one prime $p > 4\sqrt{M}$ and then to the first power. For each prime $4\sqrt{M} < p \leq M$, select some m_p for which $p|u + m_p v$. We let $\mathcal{M} = \{m_p : 4\sqrt{M} < p \leq M\}$ and apply Lemma 1, so that $|B| \geq \pi(M) - \pi(4\sqrt{M})$.

We can do better than this. We can take

$$\mathcal{M} = \{m_p : 4\sqrt{M} < p \leq M\} \cup \{p > M : p = u + mv \text{ for some } 1 \leq m \leq M\}$$

so that $|\mathcal{M}| = \pi(M) - \pi(4\sqrt{M}) + \pi(u + Mv; v, u) - \pi(\max\{M, u\}; v, u)$. This latter quantity $\pi(u + Mv; v, u) - \pi(\max\{M, u\}; v, u)$ is $> \epsilon M/\log M$ unless $v = 1$

²The absolute constants are probably not optimal, but it doesn't affect the later steps of the proof.

and $u \ll \epsilon M$. In fact to get as small as the set already constructed we must have $u \ll M^{2/3} \log M$ and $v = 1$, which we will assume from now on. $\square$

We proceed with the following simple lemma which says that the regime $v = 1, u \ll M$ is essentially the same as $v = 1, u = 0$.

Lemma 3. *Let I be an integer interval of size M contained in the interval $[2M]$. Then for any $x \in [M]$ there is $k \geq 0$ such that $2^k x \in I$.*

Proof. Let $x \in [M]$ and $I = [a+1, a+M]$ for some $0 \leq a \leq M$. If $x \in I$ we simply take $k = 0$. Otherwise, let k be the smallest integer such that $2^{k+1}x > a + M$. Then $2^k x \leq a + M$ by construction. On the other hand $2^k x > a$ since $a \leq M$, so $2^k x \in I$. $\square$

Let $\mathcal{P}_1$ be the set of primes in the interval $[M^{1/3} + 1, M]$ and $\mathcal{P}_2$ be the set of primes in the interval $[3, M^{1/3}]$. For any integer $x \in [M]$ by Lemma 3 we can pick some m_x such that $u + m_x = 2^{k(x)}x$. We define

$$\mathcal{M}_1 = \{1 \leq m_p \leq M, p \in \mathcal{P}_1\}$$

and

$$\mathcal{M}_2 = \{1 \leq m_x \leq M, x = p_i p_j p_k\},$$

where p_i, p_j, p_k are distinct primes in $\mathcal{P}_2$.

By the Prime Number Theorem we have

$$|\mathcal{M}_1| \geq \pi(M) - \pi(M^{1/3})$$

and

$$|\mathcal{M}_2| \geq \pi(M^{1/3})^3 \geq M^{1-o(1)}.$$

Obviously, $\mathcal{M}_1$ and $\mathcal{M}_2$ as well as $\mathcal{P}_1$ and $\mathcal{P}_2$ are disjoint. We will now use a map similar to the one used in the proof of Lemma 1, namely $\rho : \mathbb{Z} \rightarrow \mathbb{F}_3^{|\mathcal{P}_1|+|\mathcal{P}_2|}$. The map is defined as follows. Order the primes in $\mathcal{P}_1 \cup \mathcal{P}_2$, so that the elements in $\mathcal{P}_1$ receive a smaller index than the ones from $\mathcal{P}_2$. Then $\rho(x)_i$ (the i th coordinate of $\rho(x)$) equals to $v_{p_i}(x) \bmod 3$, where p_i is the prime in $\mathcal{P}_1 \cup \mathcal{P}_2$ with index i . Consider the set $B' = \rho(B) - \rho(g)/2$. Then, $\rho(u + m) \in B' + B'$, in particular for $m \in \mathcal{M}_1 \cup \mathcal{M}_2$. Let $A' = \rho(u + m)$ with m restricted to $\mathcal{M}_1 \cup \mathcal{M}_2$.

Now we are going to look closely at what A' looks like. It will be convenient to introduce the following definition.

Definition 1. *The k -sphere in $\mathbb{F}_q^n$ is the set $S_k \subset \mathbb{F}_q^n$ of $0-1$ vectors with exactly k non-zero coordinates.*

Then our construction of $\mathcal{M}_1$ and $\mathcal{M}_2$ reveals that A' restricted to the first $|\mathcal{P}_1|$ coordinates is a 1-sphere, and restricted to the last $|\mathcal{P}_2|$ coordinates is a 3-sphere.

We now state a lemma about 3-spheres, which might be of independent interest, but defer the proof till the end of the current discussion.

Lemma 4. *Let S_3 be the 3-sphere in $\mathbb{F}_3^n$. If $S_3 \subset B + B$ then $|B| = \Omega(n^2)$.*

Using Lemma 4 as a black box, we now finish the proof of Theorem 1.

Proof of Theorem 1. Let $A' \subset B' + B'$ be the sets defined in the preceding discussion. Let H be a graph with B' as the vertex set and for each $a \in A'$ a single edge is placed joining some pair (b, b') with $b + b' = a$. In fact, the edges are of two types: the ones corresponding to $m \in \mathcal{M}_1$ and to $m \in \mathcal{M}_2$. Consider the subgraph H' spanned by the edges of the first type, so $E(H')$ restricted to the first $|\mathcal{P}_1|$ coordinates is a 1-sphere and the projection to the last $|\mathcal{P}_2|$ coordinates is zero.

Let $H_1, \dots, H_K$ be the connected components of H' . It follows that $V(H_i)|_{\mathcal{P}_2}$ (i.e. projection to the last $|\mathcal{P}_2|$ coordinates) can attain only two values on each component and they must sum to zero. Denote these vectors by $\vec{v}_i$ and $-\vec{v}_i$. Moreover, if H_i contains an odd cycle, $\vec{v}_i = \vec{0}$, since chasing such a cycle implies $\vec{v}_i = -\vec{v}_i$. On the other hand, H_i cannot contain even cycles at all, since it would imply a linear dependence between edges, which is absurd as they form a basis for the subspace spanned by the first $|\mathcal{P}_1|$ coordinates.

Let T be the number of trees among H_i . Summing up, we have the following observations. First, $2T + 1 \geq |V(H')|_{\mathcal{P}_2}|$, since any component which is not a tree has trivial projection and a tree contributes at most two vectors. Second, $|V(H_i)| \geq |E(H_i)|$ since $V(H_i)$ must span the subspace generated by $E(H_i)$ which has dimension $|E(H_i)|$. Of course, in the case of a tree we have $|V(H_i)| = |E(H_i)| + 1$. Thus, we have

$$|\mathcal{M}_1| + T = \sum_i |E(H_i)| + T \leq |V(H')|.$$

Let now $B_S = (B' \setminus V(H'))|_{\mathcal{P}_2} \cup V(H')|_{\mathcal{P}_2}$. It is trivial that $B_S|_{\mathcal{P}_2} = B'|_{\mathcal{P}_2}$. Thus, $B_S|_{\mathcal{P}_2} + B_S|_{\mathcal{P}_2}$ contains a 3-sphere (which corresponds to the elements in $\mathcal{M}_2$), so we can apply Lemma 4 and obtain

$$|B_S| \geq |B_S|_{\mathcal{P}_2} \gg |\mathcal{P}_2|^2 \geq |\mathcal{M}_2|^{2/3-o(1)},$$

which means that at least (moving the implicit constant in $\gg$ into the $o(1)$ term)

$$|\mathcal{M}_2|^{2/3-o(1)} \leq |V(H')|_{\mathcal{P}_2} \leq 2T + 1$$

or

$$|\mathcal{M}_2|^{2/3-o(1)} \leq |(B' \setminus V(H'))_{\mathcal{P}_2}| \leq |B' \setminus V(H')|.$$

In any case we have (suppressing constants into the $o(1)$ term)

$$|\mathcal{M}_1| + |\mathcal{M}_2|^{2/3-o(1)} \leq |V(H')| + |B' \setminus V(H')| = |B'|,$$

so unwinding the definitions and using the trivial $|B'| \leq |B|$, we obtain (again hiding lower order terms in $o(1)$)

$$\pi(M) + M^{2/3-o(1)} \leq \pi(M) - \pi(M^{1/3}) + M^{2/3-o(1)} \leq |\mathcal{M}_1| + |\mathcal{M}_2|^{2/3-o(1)} \leq |B|.$$

□

It remains to prove Lemma 4.

Proof of Lemma 4. Let us examine what a difference between two elements $a, a' \in S_3$ can look like. To this end, let us fix $k \in \mathbb{F}_3^n$ and look at the number of solutions to

$$k = a - a', \quad a, a' \in S_3. \quad (3)$$

For $k \neq 0$ there are only three possibilities.

Case 1. k has three one-coordinates and three two-coordinates. Then we can uniquely reconstruct a and a' .

Case 2. k has four non-zero coordinates: two ones and two minus ones (note that in $\mathbb{F}_3$ we have $-1 = 2$). Then there are less than n solutions to (3).

Case 3. k has only two non-zero coordinates: one of them is one and the other is equal to two. Then there are less than n^2 solutions to (3).

We will say that a pair of elements (x, y) in $\mathbb{F}_3^n \times \mathbb{F}_3^n$ (not necessarily in $S_3 \times S_3$) is a Case 1, 2, 3 pair if $x - y$ has respectively 6, 4, 2 non-zero coordinates, as described above.

Let H be a graph on B such that for each $a \in S_3$ there is an edge between a single pair (b, b') for which $a = b + b'$, similar to the one we have already encountered in the proof of Theorem 1. It will be convenient to assume now that H is bipartite with two copies of B (say, B_1 and B_2) as color classes, which we can do without loss of generality³. The crucial observation is that if $v_1, v_2 \in B_1$ have a common neighbour, then the pair (v_1, v_2) is either a Case 1, 2 or 3 pair. Indeed, then there is $w \in B_2$ such that both $v_1 + w \in S_3$ and $v_2 + w \in S_3$, so there are $a, a' \in S_3$ such that $v_1 - v_2 = a - a'$.

In fact, we are going to estimate

$$\sum_{(v_1, v_2) \in B_1 \times B_1} |N(v_1) \cap N(v_2)| = \sum_{w \in B_2} d^2(w). \quad (4)$$

For convenience, we will assume henceforth that vertices denoted by the letter v belong to the first color class B_1 while those denoted by w belong to the color class B_2 . The sum in (4) naturally splits into a sum over Case 1, Case 2 and Case 3 pairs, which we will estimate separately. Our aim is to bound each contribution from above by $O(n^4)$.

The contribution given by the Case 2 pairs is the hardest one. In fact the contribution can be as large as n^5 , for example if $B_1 = B_2 = S_1 \cup S_2$. To overcome this difficulty we will excise some edges from H while still retaining a positive

³If (b_1, b_2) was an edge in the original graph with a single copy of B as the vertex set, then we place an edge between $\min(b_1, b_2)$ and $\max(b_1, b_2)$, so that the former vertex is in B_1 and the latter in B_2 .

proportion of S_3 represented by edges, but significantly reducing the contribution from Case 2 pairs.

Case 1 pairs.

For any Case 1 pair (v_1, v_2) holds $|N(v_1) \cap N(v_2)| \leq 1$ since for a common neighbour w of v_1, v_2 we can uniquely reconstruct $v_1 + w$ and $v_2 + w$. Thus, we can simply bound the whole contribution from Case 1 pairs by $|B|^2$, which we may assume is at most n^4 , otherwise we are done.

Case 3 pairs.

Now we estimate the contribution of the pairs with $b - b' = (\dots, 0, 1, \dots, 2, \dots)$. We claim that for a fixed $k = (\dots, 0, 1, \dots, 2, \dots)$ holds

$$\sum_{b-b'=k} |N(b) \cap N(b')| \leq n^2. \quad (5)$$

Let $\pi_i : \mathbb{F}_3^n \rightarrow \mathbb{F}_3$ be the projection to the i th coordinate and i_1, i_2 be the indices such that $\pi_{i_1}(k) = 1$ and $\pi_{i_2}(k) = 2$, respectively. For an arbitrary common neighbor b_c of b and b' holds

$$(b + b_c) - (b' + b_c) = b - b' = k.$$

Thus, $\pi_{i_1}(b + b_c) = \pi_{i_2}(b' + b_c) = 1$ and, moreover, there are exactly two indices j_1, j_2 such that

$$\pi_{j_1, j_2}(b + b_c) = \pi_{j_1, j_2}(b' + b_c) = 1$$

and other coordinates are zero. We can thus map b_c to the vector v with $\pi_{j_1, j_2}(v) = 1$. The set of common neighbours of Case-3 pairs (b, b') with $b - b' = k$

$$\bigcup_{b-b'=k} (N(b) \cap N(b'))$$

is then mapped the set of vectors $v \in \mathbb{F}_3^n$ for which $\pi_{j_1}(v) = \pi_{j_2}(v) = 1$ for some $j_1 \neq j_2$ with the rest of coordinates being zero. The map is in fact injective, since we can reconstruct $b + b_c$ and $b' + b_c$ from k and v . In turn, the pair of edges corresponding to $b + b_c$ and $b' + b_c$ uniquely identifies the pair (b, b') . Obviously the number of v 's is bounded by n^2 , so we get (5).

Finally, there are at most n^2 choices of k in (5), so the total contribution from Case 3 pairs to (4) is $O(n^4)$.

Case 2 pairs.

Let us get back to Case 2 pairs. The graph H can be naturally decomposed into the union of graphs H_i , $i = 1 \dots n$, where $E(H_i) = \bar{S}_i := \{v \in S_3 | \pi_i(v) = 1\}$ and $V(H_i)$ is the set of vertices spanned by $E(H_i)$. In fact, $\bar{S}_i$ is a shifted 2-sphere,

so this decomposition corresponds to the decomposition of S_3 into a union of n shifted two-spheres given by fixing the i th coordinate to one.

Let $W_i \subset V(H_i)|_{B_2}$ be the set of vertices w of H_i in B_2 such that $d_{H_i}(w) > 2^{10}n$, where d_{H_i} denotes degree in H_i . We claim that

$$\sum_{w \in W_i} d_{H_i}(w) \leq \frac{n^2}{50}. \quad (6)$$

The claim follows from the sublemma below which essentially asserts that a large portion of the 2-sphere cannot be additively split into two unequal parts unless the size of one of the parts is comparable to the size of the whole sphere.

Lemma 5. *Let $X, Y \subset \mathbb{F}_3^n$ with $|X| \leq n/2^{10}$ and $|Y| \leq n^2/100$. Then*

$$|(X + Y) \cap S_2| \leq \frac{n^2}{50}.$$

Before we proceed with the proof, let us show how the claimed bound (6) follows from Lemma 5. Clearly,

$$|W_i| \leq \frac{|E(H_i)|}{2^{10}n} \leq \frac{n}{2^{10}}.$$

Let $\bar{S}'_i$ be the subset of $\bar{S}_i$ represented by all edges emanating from W_i . We have that $\bar{S}'_i \subset B_1 + W_i$. Equivalently, $\bar{S}'_i - 1_i \subset (B_1 - 1_i) + W_i$, where 1_i is the vector with $\pi_i(1_i) = 1$ and zero coordinates otherwise. Now either $|B_1| \geq n^2/100$ and we are done with the whole Lemma 4, or we apply Lemma 5 with $X = W_i$ and $Y = B_1 - 1_i$, so that

$$|\bar{S}'_i| \leq |(X + Y) \cap S_2| \leq \frac{n^2}{50}.$$

But $|\bar{S}'_i| = \sum_{w \in W_i} d_i(w)$ by construction.

Proof of Lemma 5. The intuition behind the proof is as follows. Let L be the subspace spanned by X , so its dimension is at most $|X| \ll n$. Let π be the projection to $L^\perp$ and assume that $|\pi(S_2)| \approx |S_2|$ (we don't specify the exact meaning of $\approx$ here, but since L is small, one may expect that $\pi(S_2)$ is comparable in size to the whole sphere). Then we can estimate

$$|S_2 \setminus ((X + Y) \cap S_2)| \geq |\pi(S_2 \setminus ((X + Y) \cap S_2))| \approx |S_2| - |Y|,$$

since $|\pi(X + Y)| = |\pi(Y)| \leq |Y|$. Thus, $|(X + Y) \cap S_2| \approx |Y|$.

Now we implement the outlined strategy. Let L be subspace of $\mathbb{F}_3^n$ spanned by X , $l := \dim L$ and $m := n - l$, so $m \geq (1 - 2^{-10})n$. Let $e_1, \dots, e_m$ be standard basis elements, so $e_i = 1_i$ for $i = 1, \dots, m$. Without loss of generality we may assume that $\{L, e_1, \dots, e_m\}$ spans the whole space $\mathbb{F}_3^n$. Then let T be a linear invertible transformation such that $T(e_i) = e_i$ for $1 \leq i \leq m$ and $T(L) = \text{Span}(\{e_{m+1}, \dots, e_n\})$. Denote $M := \text{Span}(\{e_1, \dots, e_m\})$, so in particular $T(X) \perp M$.

We have (as T is one-to-one)

$$\begin{aligned} |(X + Y) \cap S_2| &= |T(X + Y) \cap T(S_2)| \\ &= |T(X + Y) \cap T(S_2) \cap M| + |T(X + Y) \cap T(S_2) \cap M^c|. \end{aligned} \quad (7)$$

Since $T(X)$ has trivial orthogonal projection to M and $T(X + Y) = T(X) + T(Y)$, we can crudely estimate

$$\begin{aligned} |T(X + Y) \cap T(S_2) \cap M| &\leq |T(X + Y) \cap M| \\ &\leq |\pi_M(T(X + Y))| = |\pi_M(T(Y))| \leq |Y|, \end{aligned} \quad (8)$$

where π_M denotes orthogonal projection onto M . For the second term we have

$$|T(X + Y) \cap T(S_2) \cap M^c| \leq |T(S_2) \cap M^c| = |T(S_2)| - |T(S_2) \cap M|. \quad (9)$$

But since $T(e_i + e_j) = T(e_i) + T(e_j) = e_i + e_j \in M$ for $1 \leq i, j \leq m$,

$$\begin{aligned} |T(S_2) \cap M| &\geq \frac{m(m-1)}{2} \geq \frac{((1-2^{-10})n)((1-2^{-10})n-1)}{2} \\ &\geq 0.991 \frac{n(n-1)}{2} \geq 0.99 \frac{n^2}{2}, \end{aligned}$$

for n large enough. Therefore,

$$|T(X + Y) \cap T(S_2) \cap M^c| \leq |T(S_2)| - 0.99 \frac{n^2}{2} \leq 0.01 \frac{n^2}{2}. \quad (10)$$

Combining (7), (8) and (10) we get

$$|(X + Y) \cap S_2| \leq 0.01 \frac{n^2}{2} + \frac{n^2}{100} \leq \frac{n^2}{50},$$

as required. □

We proceed with the proof of Lemma 4.

We prune each H_i , cutting all edges emanating from each $w \in W_i$, that is, from those vertices whose degree in H_i is at least $2^{10}n$. By (6), we will remove at most $n^2/50$ edges in total. Let us call the pruned graph H'_i and count the number of common neighbours of all Case 2 pairs in H'_i . To emphasize the fact that we are counting common neighbours in H'_i now, we denote by $N_{H'_i}(v)$ the set of neighbours of a vertex v in H'_i . Note, however, that the property that a pair of vertices (v_1, v_2) is a Case 2 pair is independent of the graph they belong to.

We have

$$\sum_{(v_1, v_2) \text{ Case 2 in } H'_i} |N_{H'_i}(v_1) \cap N_{H'_i}(v_2)| \leq \sum_{w \in V(H'_i) \setminus B_2} d_{H'_i}^2(w).$$

On the other hand, by construction of H'_i , we have

$$\begin{aligned} \sum_{w \in V(H'_i)|_{B_2}} d_{H'_i}^2(w) &\leq \max_{w \in V(H'_i)|_{B_2}} d_{H'_i}(w) \left(\sum_{w \in V(H'_i)|_{B_2}} d_{H'_i}(w) \right) \\ &\leq 2^{10} n |\bar{S}_i| \ll n^3. \end{aligned} \quad (11)$$

Let $H' := \bigcup_i H'_i$. By (6),

$$|E(H')| \geq |E(H)| - \frac{n^3}{50} \gg n^3. \quad (12)$$

Also, we claim that if $v_1 w, v_2 w \in E(H')$ and (v_1, v_2) is a Case 2 pair in H' , then there is an i such that $v_1 w, v_2 w \in E(H'_i)$, that is, (v_1, v_2) is a Case 2 pair in H'_i . Indeed, there is an i such that $\pi_i(v_1 + w) = \pi_i(v_2 + w) = 1$, but then $v_1 w, v_2 w \in E(H'_i)$ by construction. Thus, by (11),

$$\sum_{(v_1, v_2) \text{ Case 2 in } H'} |N_{H'}(v_1) \cap N_{H'}(v_2)| \leq \sum_{i=1}^n \sum_{(v_1, v_2) \text{ Case 2 in } H'_i} |N_{H'_i}(v_1) \cap N_{H'_i}(v_2)| \ll n^4. \quad (13)$$

Now for the graph H' we combine the estimates for Case 1, Case 2 and Case 3 pairs (as $E(H') \subset E(H)$ we can safely use the upper bounds obtained for H in Cases 1 and 3) and conclude

$$\begin{aligned} \sum_{w \in B_2} d_{H'}^2(w) &= \sum_{(v_1, v_2) \in B_1 \times B_1} |N_{H'}(v_1) \cap N_{H'}(v_2)| \\ &= \sum_{v_1 - v_2 \text{ Case 1}} |N_{H'}(v_1) \cap N_{H'}(v_2)| \\ &\quad + \sum_{v_1 - v_2 \text{ Case 2}} |N_{H'}(v_1) \cap N_{H'}(v_2)| \\ &\quad + \sum_{v_1 - v_2 \text{ Case 3}} |N_{H'}(v_1) \cap N_{H'}(v_2)| \\ &\ll n^4. \end{aligned} \quad (14)$$

By the Cauchy-Schwartz inequality

$$\left(\sum_{w \in B_2} d_{H'}(w) \right)^2 \leq |B_2| \left(\sum_{w \in B_2} d_{H'}^2(w) \right),$$

so (12) and (14) give

$$n^6 \ll |E(H')|^2 \ll \left(\sum_{w \in B_2} d_{H'}(w) \right)^2 \ll |B_2| n^4.$$

Therefore, $|B_2| = |B| = \Omega(n^2)$.

□

3. ACKNOWLEDGMENTS

The author is grateful to Andrew Granville for various refinements of the original argument, in particular for optimizing the lower bound, and for bringing the problem back to the author's attention. The author would also like to thank Peter Hegarty for very careful proofreading and Ilya Shkredov for discussions and coining the term “discrete sphere”.

REFERENCES

- [1] E. Croot and V. Lev, Open problems in additive combinatorics. Additive combinatorics, 207–233, CRM Proc. Lecture Notes, 43, Amer. Math. Soc., Providence, RI, 2007.
- [2] O. Roche-Newton and D. Zhelezov, A bound on the multiplicative energy of a sum set and extremal sum-product problems, *Moscow J. Comb. Number Theory* 5 (2015), no. 1.
- [3] T. Tao, The sum-product phenomenon in arbitrary rings. *Contrib. Discrete Math.* 4 (2009), no. 2, 59–82.
- [4] T. Tao and V. Vu, Additive combinatorics, Cambridge Studies in Mathematics, Cambridge Univ. Press, 2006.
- [5] D. Zhelezov, On sets with small additive doubling in product sets, to appear in *J. Number Theory*.
- [6] D. Zhelezov, Improved bounds for arithmetic progressions in product sets, to appear in *Int. J. Number Theory*.